

1 XOR

Note 1

The truth table of XOR (denoted by \oplus) is as follows.

A	B	$A \oplus B$
F	F	F
F	T	T
T	F	T
T	T	F

(a) Express XOR using only (\wedge, \vee, \neg) and parentheses.

(b) Does $(A \oplus B)$ imply $(A \vee B)$? Explain briefly.

(c) Does $(A \vee B)$ imply $(A \oplus B)$? Explain briefly.

2 Proof Practice

Note 2

(a) Prove that $\forall n \in \mathbb{N}$, if n is odd, then $n^2 + 1$ is even. (Recall that n is odd if $n = 2k + 1$ for some natural number k .)

(b) Prove that $\forall x, y \in \mathbb{R}, \min(x, y) = (x + y - |x - y|)/2$. (Recall, that the definition of absolute value for a real number z , is

$$|z| = \begin{cases} z, & z \geq 0 \\ -z, & z < 0 \end{cases}$$

(c) Suppose $A \subseteq B$. Prove $\mathcal{P}(A) \subseteq \mathcal{P}(B)$. (Recall that $A' \in \mathcal{P}(A)$ if and only if $A' \subseteq A$.)

3 Numbers of Friends

Note 2 Prove that if there are $n \geq 2$ people at a party, then at least 2 of them have the same number of friends at the party. Assume that friendships are always reciprocated: that is, if Alice is friends with Bob, then Bob is also friends with Alice.

(Hint: The Pigeonhole Principle states that if n items are placed in m containers, where $n > m$, at least one container must contain more than one item. You may use this without proof.)

4 Preserving Set Operations

Note 0
Note 2

For a function f , define the image of a set X to be the set $f(X) = \{y \mid y = f(x) \text{ for some } x \in X\}$. Define the inverse image or preimage of a set Y to be the set $f^{-1}(Y) = \{x \mid f(x) \in Y\}$. Prove the following statements, in which A and B are sets.

Recall: For sets X and Y , $X = Y$ if and only if $X \subseteq Y$ and $Y \subseteq X$. To prove that $X \subseteq Y$, it is sufficient to show that $(\forall x) ((x \in X) \implies (x \in Y))$.

(a) $f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$.

(b) $f(A \cup B) = f(A) \cup f(B)$.